

Genomic Landscape of Waldenström Macroglobulinemia and Its Impact on Treatment Strategies

Steven P. Treon, MD, PhD^{1,2}; Lian Xu, MS¹; Maria Luisa Guerrero, MD^{1,2}; Cristina Jimenez, PhD^{1,2}; Zachary R. Hunter, PhD^{1,2}; Xia Liu, MD^{1,2}; Maria Demos, BS¹; Joshua Gustine, MPH¹; Gloria Chan, MS¹; Mani Munshi, MS¹; Nicholas Tsakmaklis, BA¹; Jiaji G. Chen, BS¹; Amanda Kofides, BS¹; Romanos Sklavonitis-Pistofidis, MD^{2,3,4}; Mark Bustoros, MD^{2,3}; Andrew Keezer, BS¹; Kirsten Meid, MPH¹; Christopher J. Patterson, MAcc, MPH¹; Antonio Sacco, RN^{3,4}; Aldo Roccaro, MD, PhD⁴; Andrew R. Branagan, MD⁵; Guang Yang, PhD^{1,2}; Irene M. Ghobrial, MD^{2,3}; and Jorge J. Castillo, MD^{1,2}

Next-generation sequencing has revealed recurring somatic mutations in Waldenström macroglobulinemia (WM), including *MYD88* (95%-97%), *CXCR4* (30%-40%), *ARID1A* (17%), and *CD79B* (8%-15%). Deletions involving chromosome 6q are common in patients with mutated *MYD88* and include genes that modulate NFκB, BCL2, Bruton tyrosine kinase (BTK), and apoptosis. Patients with wild-type *MYD88* WM show an increased risk of transformation and death and exhibit many mutations found in diffuse large B-cell lymphoma. The discovery of *MYD88* and *CXCR4* mutations in WM has facilitated rational drug development, including the development of BTK and CXCR4 inhibitors. Responses to many agents commonly used to treat WM, including the BTK inhibitor ibrutinib, are affected by *MYD88* and/or *CXCR4* mutation status. The mutation status of both *MYD88* and *CXCR4* can be used for a precision-guided treatment approach to WM.

J Clin Oncol 38:1198-1208. © 2020 by American Society of Clinical Oncology

INTRODUCTION

Waldenström macroglobulinemia (WM) is a B-cell malignancy characterized as an immunoglobulin M (IgM)-secreting lymphoplasmacytic lymphoma using WHO criteria.¹ Patients can present with morbidity related to excess malignant lymphoplasmacytic cells (LPCs) in the bone marrow (BM), lymph nodes, and spleen, as well as IgM production that can produce symptomatic hyperviscosity, cryoglobulinemia, and autoimmune-related complications.² Despite advances in therapy, WM remains incurable. Treatment options for WM have largely represented therapeutics studied in other diseases, with more recent advances facilitated by next-generation sequencing (NGS).

Using NGS, recurring somatic mutations in *MYD88*, *CXCR4*, *ARID1A*, and *CD79* were identified in WM LPC. Copy number alterations, including those in chromosome 6q that affect regulatory genes for NFκB, Bruton tyrosine kinase (BTK), BCL2, and apoptotic signaling, were also identified.³ Although most patients with WM (95%-97%) carry a point mutation in *MYD88* that switches leucine to proline at amino acid position 265, those that are wild type for *MYD88* show a more aggressive disease course and possess somatic mutations that overlap with those found in diffuse large B-cell lymphoma (DLBCL).^{4,5} Herein, we discuss the genomic landscape of WM and the impact of underlying genomics on disease presentation, transcriptional changes, treatment outcome, and overall survival. The use of *MYD88* and *CXCR4* mutation status to guide treatment

in treatment-naïve and previously treated patients with WM is also discussed.

MUTATIONS IN *MYD88*

A recurring somatic mutation in *MYD88* (L265P) was identified in 91% of patients with WM by paired tumor and normal whole-genome sequencing and subsequently confirmed by Sanger sequencing and allele-specific polymerase chain reaction (AS-PCR) assays by multiple investigators.⁶ Using sensitive AS-PCR testing, *MYD88* L265P was found to be expressed in 93%-97% of patients with WM and was identified in both sorted B cells and plasma cells that make up the malignant clone in WM.⁷⁻¹¹ Non-L265P *MYD88* mutations have also been identified, although expression estimates for these variants are 1%-2% in WM.^{12,13} Mutated *MYD88* was also detectable in patients with IgM but not immunoglobulin G or immunoglobulin A monoclonal gammopathy of unknown significance (MGUS), suggesting an early oncogenic role for *MYD88* in WM pathogenesis.^{7,8,10} Patients with IgM MGUS with detectable mutated *MYD88* and patients with a higher mutated allele burden are at greater risk of progression to WM.^{10,14}

The *MYD88* L265P mutation can also be detected by AS-PCR in peripheral-blood (PB) samples, particularly in treatment-naïve patients with WM.¹⁵ Prior therapy with B-cell-depleting agents can greatly decrease detection of *MYD88* L265P in PB samples. *MYD88* L265P can also be found in skin lesions, CSF, and

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on November 25, 2019 and published at ascopubs.org/journal/jco on February 21, 2020; DOI <https://doi.org/10.1200/JCO.19.02314>

pleural effusions in patients with WM, providing a means of demonstrating extramedullary disease involvement.¹⁶⁻¹⁸ Cell-free DNA has also been used to detect *MYD88* L265P from PB samples of patients with WM and may represent a novel means for establishing *MYD88* mutation status.^{19,20}

Structural alterations on chromosome 3p can increase the allele burden of mutated *MYD88*.^{6,21} Up to 15% of untreated and 25% of previously treated patients may be homozygous for *MYD88* as a result of deletions of the wild-type *MYD88* allele, amplifications of the mutant *MYD88* allele, and, more commonly, acquired uniparental disomy (aUPD) events.²² aUPD events resulting in homozygous *MYD88* expression are associated with concurrent *CXCR4* mutations, the significance of which remains to be clarified but may be related to disease length and prior ibrutinib exposure.^{6,23}

Patients with wild-type *MYD88* show similar histologic findings and transcription profile as patients with *MYD88*-mutated WM.^{4,5} However, patients with wild-type *MYD88* with asymptomatic WM have a higher risk of symptomatic progression,²⁴ and those presenting with symptomatic disease have a greater risk of disease transformation^{8,25} and decreased overall survival.⁸ Patients with wild-type *MYD88* also show poor response to ibrutinib (discussed later). It is important to distinguish patients with suspected WM with wild-type *MYD88* disease from patients with other IgM-secreting malignancies. In one series, 30% of patients with suspected wild-type *MYD88* WM had an alternative diagnosis, including IgM multiple myeloma (MM). The presence of very high serum IgM levels, lytic lesions, and/or renal dysfunction may help identify IgM MM.^{8,26,27} Use of cytogenetics to evaluate for t(11;14) and cyclin D1 staining can be helpful in distinguishing IgM MM from wild-type *MYD88* WM.^{8,27,28}

MYD88 is an adaptor protein that interacts with the Toll-like and interleukin (IL)-1 receptors and dimerizes upon receptor activation. Dimerization of *MYD88* provides a scaffold for recruitment of other proteins to a Myddosome complex that triggers downstream signaling, leading to nuclear factor- κ B (NFKB) activation (Fig 1).²⁹ Both IRAK1/IRAK4 and BTK are Myddosome components and trigger NFKB.^{30,31} Recruitment and activation of the IRAK and BTK molecules can be blocked by either knockdown or inhibition of *MYD88* that leads to apoptosis of mutated *MYD88* WM cells. Mutated *MYD88* can also upregulate transcription of the SRC family member HCK that normally is downregulated in late stages of B-cell ontogeny and can transactivate HCK via IL-6.³² Activated HCK triggers pro-survival signaling of mutated *MYD88* WM cells through BTK, PI3K/AKT, and MAPK/ERK1/2. Both BTK and HCK are targets of ibrutinib, which has shown remarkable activity in patients with mutated *MYD88* WM.^{32,33}

More recently, a role for mutated *MYD88* as a trigger for B-cell receptor (BCR) pathway cross talk has been

proposed. Phelan et al³⁴ described the existence of a *MYD88*-TLR9-BCR(IgM) (MY-T-BCR) supercomplex in mutated *MYD88* tumor cells. Evidence for chronic activated BCR signaling exists in WM.^{35,36} SYK, a component of the BCR pathway, is present in its active form in the Myddosome, and its activation is triggered by mutated *MYD88*.³⁷ Importantly, mutated *MYD88*-directed SYK activation triggers STAT3 and AKT pro-survival signaling. In preclinical studies, the addition of an SYK inhibitor triggered synthetic lethality when combined with ibrutinib.³⁷ Taken together, these studies show that mutated *MYD88* can direct multiple pro-survival cascades that lead to NFKB, AKT, ERK, and STAT3 activation in WM cells (Fig 1). WM cells also export mutated *MYD88* via extracellular vesicles, which may produce a growth-supportive proinflammatory microenvironment (Fig 1).³⁸

MUTATIONS IN *CXCR4*

Somatic mutations involving the C-terminal domain of *CXCR4* are present in up to 40% of patients with WM, and although they almost always occur in those with *MYD88* mutations, some patients with wild-type *MYD88* can also express *CXCR4* mutations.^{21,39} *CXCR4* mutations are essentially unique to WM, with only a few cases of marginal zone lymphoma and activated B-cell (ABC) subtype of DLBCL reported so far. Germline mutations in the C-terminal domain of *CXCR4* are present in patients with WHIM (autosomal dominant warts, hypogammaglobulinemia, infection, and myelokathexis) syndrome.⁴⁰ In patients with WHIM syndrome, activation of *CXCR4* by its ligand CXCL12 induces extended chemotactic signaling and neutrophil sequestration in the BM (myelokathexis) and impairment of lymphocyte development. In WM, > 40 nonsense and frameshift mutations in the C-terminal domain of *CXCR4* have been described.^{41,42} Mutations in the C-terminal domain of *CXCR4* lead to loss of regulatory serines and promote sustained CXCL12-mediated activation of the AKT and ERK pathways and in vivo WM disease progression and dissemination in mice.⁴³⁻⁴⁵ Despite the autonomous pro-survival signaling associated with *CXCR4* mutations, inhibition of *MYD88* triggers apoptosis in both wild-type and mutated *CXCR4*-expressing WM cells, consistent with a primary role for mutated *MYD88* survival signaling in WM.⁴⁵

Unlike *MYD88*, mutated *CXCR4* is subclonal, with highly variable clonality averaging approximately 35%. Multiple *CXCR4* mutations can also exist within individual patients that can occur in separate clones or as compound heterozygous events.³⁹ The subclonal nature of *CXCR4* mutations relative to *MYD88* and their paucity in IgM MGUS relative to WM suggest that *CXCR4* mutations are acquired after *MYD88*.³⁹ Clonal 6q deletions (del 6q21-25), which are found in 40%-50% of patients with WM, appear exclusive of *CXCR4* in treatment-naïve patients.⁴⁶ The increasing frequency of 6q deletions from IgM MGUS




FIG 1. Prosurvival signaling mediated by mutated *MYD88*. Mutated *MYD88* triggers assembly of the Myddosome, which includes activated BTK and IRAK4/IRAK1 that transactivate NFκB. Mutated *MYD88* also transcriptionally upregulates and transactivates through interleukin (IL)-6 the SRC family member HCK, which triggers activation of BTK itself, as well as AKT and ERK. Mutated *MYD88* also mediates cross talk through LYN-activated SYK, which activates STAT3 and AKT prosurvival signaling. Deletions in chromosome 6q result in the loss of important regulators of *MYD88* signaling, including inhibitor of BTK (IBTK) and the NFκB regulators HIVEP2 and TNFAIP3. Waldenström macroglobulinemia cells also export mutated *MYD88* via extracellular vesicles to induce signaling cascades in mast cells and macrophages that provide a supportive proinflammatory microenvironment. BCR, B-cell receptor; mTOR, mammalian target of rapamycin; TLRs, Toll-like receptors.

through asymptomatic and symptomatic WM suggests that loss of genes within this region (discussed later) facilitates disease progression.^{13,35,46} Taken together, these findings suggest that at least 2 distinct pathways mediated by either mutated *CXCR4* or del 6q21-25 promote IgM MGUS progression to symptomatic WM.

Like *MYD88*, *CXCR4* somatic mutations can affect WM disease presentation. Patients with mutated *CXCR4* present with less adenopathy, and those with nonsense *CXCR4* mutations have increased BM disease, serum IgM levels, and/or symptomatic hyperviscosity at presentation.⁴¹ Nonsense *CXCR4* mutations are also associated with acquired

von Willebrand factor deficiency.^{41,47} Despite differences in clinical presentation, *CXCR4* mutations do not adversely affect overall survival but do seem to affect time to treatment.^{8,13,41}

In vitro modeling of WM cells transduced with mutated *CXCR4* reveals increased drug resistance in the presence of its ligand CXCL12 to multiple therapeutics including bendamustine, fludarabine, bortezomib, idelalisib, and ibrutinib.⁴³⁻⁴⁵ These studies also showed that resistance mediated by mutated *CXCR4* could be reversed by use of *CXCR4*-blocking agents in both in vitro and in vivo studies.⁴³⁻⁴⁵

OTHER RECURRING MUTATIONS

Somatic mutations in *ARID1A* are present in 17% of patients with WM, including nonsense and frameshift variants.²¹ Patients with *ARID1A* and *MYD88* L265P mutations have greater WM disease involvement and lower hemoglobin and platelet counts. *ARID1A* and its frequently deleted homolog *ARID1B* (see later discussion) are on chromosome 6q.⁴⁶ Both are switch/sucrose nonfermentable (SWI/SNF) family members and serve as chromatin remodeling genes, thereby modulating gene regulation. Although still poorly understood, *ARID1A* can modulate TP53 and is thought to act as an epigenetic tumor suppressor in ovarian cancer.³

Mutations in *CD79A* and *CD79B* can be found in 8%-12% of patients with WM.^{13,21,42} Both are components of the BCR pathway and can form heterodimers with each other.³ The CD79A/B heterodimer associates with the immunoglobulin heavy chain required for cell surface expression of BCR and BCR-induced signaling. Activating mutations in the immunotyrosine-based activation motif of CD79A and CD79B occur in the ABC subtype of DLBCL and trigger SYK, PLC γ 2, and BTK.⁴⁸ Mutated MYD88 cross talk through SYK may also explain chronic BCR signaling observed in WM cells.³⁷ Deletions of *LYN*, which are found in 70% of patients with WM, may contribute to hyperresponsive BCR signaling, as informed by *lyn*^{-/-} transgenic mice.^{21,49} Although mutations in both *CD79A* and *CD79B* are mainly found in patients with mutated *MYD88* including those with WM, a *CD79B* mutation was observed in a patient with wild-type *MYD88* WM.²¹ In one study, mutations in *CD79A* and *CD79B* were nearly exclusive of *CXCR4* mutations, suggesting that *CD79A/B* mutations may also have an independent role in facilitating mutated *MYD88*-directed progression in WM.⁴² In a small series of patients with WM, the coexpression of *CD79B* and *MYD88* mutations was associated with disease transformation.⁵⁰ Mutations in *TP53* are more rarely observed (2%-3%) in WM^{13,21} and are associated with poor survival.^{51,52}

MUTATIONS IN WILD-TYPE *MYD88* WM

A small number of patients with WM (3%-7%) lack mutations in *MYD88*, including non-L265P mutations.^{5,21} Their disease course is marked by an increased risk of disease transformation^{5,25} and shorter overall survival.⁵ Moreover, patients with wild-type *MYD88* show little response to ibrutinib (discussed further later). These findings point to fundamental differences in underlying genomics. NGS identified somatic mutations in patients with wild-type *MYD88* WM that are predicted to trigger NFKB (*TBL1XR1*, *PTPN13*, *MALT1*, *BCL10*, *NFKB1*, *NFKB2*, *NFKBIB*, *NFKBIZ*, and *UDRL1F*), impart epigenomic dysregulation (*KMT2D*, *KMT2C*, and *KDM6A*), or impair DNA damage repair (*TP53*, *ATM*, and *TRRAP*).⁵ Predicted NFKB activating mutations were downstream of BTK, and many overlapped with somatic mutations found in DLBCL.⁵ Transcriptome studies revealed a distinctive transcriptional profile in patients

with wild-type *MYD88* WM, although the most differentially expressed genes overlapped with those found in mutated *MYD88* WM.⁵ These findings may help explain many of the uniform characteristics shared among patients with WM, regardless of their underlying *MYD88* mutation status.

COPY NUMBER ALTERATIONS

Copy number alterations are common in patients with *MYD88*-mutated WM and involve both chromosome 6q and non-chromosome 6q regions.^{13,21,35} In chromosome 6q, loss of genes that modulate NFKB activity (*TNFAIP3* and *HIVEP2*), BCL2 (*BCLAF1*), apoptosis (*FOXO3*), BTK (*IBTK*), plasma cell differentiation (*PRDM1*), and *ARID1B* occur.^{21,46} Non-chromosome 6q genes that are commonly deleted include *ETV6*, a transcription repressor; *BTG1*, which is often deleted in DLBCL and associated with glucocorticoid resistance in acute lymphocytic leukemia; and *LYN*, a kinase that regulates BCR signaling.^{21,48,49} *PRDM2* and *TOP1*, which participate in TP53-related signaling, are also deleted in many patients with WM. In contrast to *MYD88*-mutated WM, recurring copy number alterations are rare in *MYD88* wild-type WM, including loss of chromosome 6q.^{5,13,21}

IMPACT OF GENOMIC ALTERATIONS ON WM TREATMENT OUTCOME

The impact of *MYD88* and *CXCR4* mutations on treatment outcome has been evaluated in several clinical trials (Table 1). Patients with *MYD88* mutations who are wild type for *CXCR4* have better outcomes associated with ibrutinib monotherapy, including time to major response, depth of response, and progression-free survival.^{33,53,54} Indeed, patients with both *MYD88* and *CXCR4* mutations exhibit a delay by 4-5 months in attaining a major response to ibrutinib monotherapy. No major responses were observed in previously treated patients with WM with wild-type *MYD88* treated with ibrutinib alone.^{12,55} Patients with *MYD88* wild-type status also had much short progression-free survival on ibrutinib alone compared with *MYD88*-mutated patients, regardless of their *CXCR4* mutation status.⁵³

Earlier disease progression has also been observed in *MYD88*- and *CXCR4*-mutated patients treated with ibrutinib alone or with ibrutinib and rituximab when compared with patients with mutated *MYD88* alone.^{53,56} Somatic mutations at the site of ibrutinib-BTK binding (Cys481) have been observed in patients with WM who experienced progression on ibrutinib.⁵⁷ Multiple *BTK* mutations were observed in this study within individual patients with WM, particularly those with *CXCR4* mutations. Although the findings highlight the importance of BTK as a treatment target in *MYD88*-mutated WM, they also raise questions as to why *CXCR4*-mutated patients are particularly at risk for the acquisition of *BTK* Cys481 mutations.

TABLE 1. Summary of the Impact of *MYD88* and *CXCR4* Mutation Status on Treatment Outcomes in Patients With Waldenström Macroglobulinemia

Agent	<i>MYD88</i>	<i>CXCR4</i>
BTK inhibitors		
Ibrutinib monotherapy	No major responses and short PFS in <i>MYD88</i> wild-type patients in prospective study	Fewer major responses and VGPRs, increased time to response, and shorter PFS in <i>CXCR4</i> -mutated patients in prospective studies
Ibrutinib plus rituximab		Fewer major responses and VGPRs and shorter PFS in <i>CXCR4</i> -mutated patients in prospective study
Zanubrutinib		Fewer major responses and VGPRs in <i>CXCR4</i> -mutated patients in prospective study (preliminary)
Proteasome inhibitors		
Bortezomib-based therapy		No impact of <i>CXCR4</i> mutation status on response rates, PFS, and OS in retrospective study
Carfilzomib, dexamethasone, rituximab		No impact of <i>CXCR4</i> mutation status on response rates, PFS, and OS in prospective study
Ixazomib, dexamethasone, rituximab		Longer median time to response in <i>CXCR4</i> -mutated patients in prospective study; response rates and PFS not affected by <i>CXCR4</i> mutation status
Alkylators		
Bendamustine-based therapy	Shorter PFS in <i>MYD88</i> wild-type patients in prospective study of bendamustine plus rituximab; no impact of <i>MYD88</i> status on overall response attainment, but PFS and TTNT were shorter in <i>MYD88</i> wild-type patients in a retrospective study	No impact of <i>CXCR4</i> mutation status on PFS with bendamustine plus rituximab in retrospective study
Cyclophosphamide, dexamethasone, rituximab	No impact of <i>MYD88</i> status on overall response attainment, but PFS and TTNT were shorter in <i>MYD88</i> wild-type patients in a retrospective study	
mTOR inhibitor		
Everolimus	Nonresponse in <i>MYD88</i> wild-type patients in a prospective study	Lower overall and major response rates in <i>CXCR4</i> -mutated patients in a prospective study
BCL2 inhibitor		
Venetoclax		Lower major response and VGPR rates in <i>CXCR4</i> -mutated patients in a prospective study (preliminary)

NOTE. Overall response rate includes minor responses or better. Major response rate includes partial responses or better. Data adapted from Owen et al.⁸⁷ Abbreviations: BTK, Bruton tyrosine kinase; mTOR, mammalian target of rapamycin; OS, overall survival; PFS, progression-free survival; TTNT, time to next treatment; VGPR, very good partial response.

The presence of *CXCR4* mutations has also affected clinical outcomes with other agents. Fewer major responses were preliminarily reported with the BTK inhibitor zanubrutinib, as well as with the BCL2 inhibitor venetoclax, in prospective studies in patients with WM.^{58,59} Time to response was also reported to be longer in patients with WM with *CXCR4* mutations receiving ixazomib-based therapy, a proteasome inhibitor (PI), in a prospective phase II study.⁶⁰ Conversely, a prospective study of patients with *CXCR4*-mutated WM treated with carfilzomib-based therapy showed no effect on clinical outcome.⁶¹ In a retrospective study, Sklavenitis-Pistofidis et al⁶² showed that bortezomib-based therapy could overcome the negative impact of *CXCR4* mutations on progression-free and overall survival.

Among treatment-naïve patients with WM receiving everolimus as a single agent in a prospective phase II study, nonresponders included patients who were *CXCR4* mutated,

as well as those who were wild type for *MYD88*.⁶³ In a retrospective study by the French Innovative Leukemia Organization group, the use of bendamustine plus rituximab resulted in similar progression-free survival among *MYD88*-mutated patients with and without *CXCR4* mutations. Conversely, those with wild-type *MYD88* showed earlier progression.⁶⁴ In a retrospective study, Paludo et al⁶⁵ also observed a trend for shorter progression-free survival and time to next treatment among genotyped patients with *MYD88* wild-type WM who received alkylator-based (bendamustine or cyclophosphamide) therapy. There were no differences in overall response rates between genotyped patients who were *MYD88* mutated or wild type, and no data on major responses were reported.

Although the previously discussed studies indicate that *CXCR4* represents an important target in WM, differences in response outcome could be modulated by the type of

CXCR4 mutation as well as its clonal presence. In a study by Castillo et al,⁶⁶ patients on ibrutinib with nonsense mutations showed fewer major responses and shorter progression-free survival compared with those with frameshift *CXCR4* mutations or those wild-type for *CXCR4*. The clonality of *CXCR4* may also be an important determinant of ibrutinib response. Gustine et al⁶⁷ investigated the impact of the *CXCR4* S338X nonsense variant on ibrutinib response outcome in patients with WM. Patients with a *CXCR4* S338X clonality of > 25% showed greater BM disease involvement and serum IgM levels, fewer very good partial responses, and shorter progression-free survival.

Given the importance of *CXCR4* mutation status in WM, a clinical trial assessing the impact of the *CXCR4* inhibitor ulocuplumab with ibrutinib in patients with *CXCR4*-mutated WM was initiated (ClinicalTrials.gov identifier: [NCT03225716](https://clinicaltrials.gov/ct2/show/study/NCT03225716)). Mavorixafor, an orally administered *CXCR4* antagonist, is also in clinical development for *CXCR4*-mutated WM. Although the use of antagonists to counter *CXCR4* signaling in *CXCR4*-mutated patients represents a rational approach for advancing treatment in this population with WM, a targeted

approach for those with *MYD88* wild-type WM remains elusive and represents an urgent priority.

GENOMIC-BASED TREATMENT APPROACH TO WM

MYD88 and *CXCR4* mutation status may be useful in treatment selection for symptomatic patients. Treatment algorithms recommended by the authors for the use of *MYD88* and *CXCR4* mutation status in the treatment approach of symptomatic untreated and previously treated patients are presented in Figures 2 and 3, respectively. In treatment-naïve patients, as well as those with previously treated WM, plasmapheresis should be considered as the first step in patients presenting with symptomatic hyperviscosity, rapidly progressing peripheral neuropathy attributed to demyelinating IgM disease, severe cryoglobulinemia, and cold agglutininemia.^{68,69} To minimize risks of short- and long-term treatment-related adverse effects associated with chemoimmunotherapy, including neuropathy, immunosuppression, and stem-cell damage, as well as secondary myelodysplasia and malignancies, the use of ibrutinib can be considered for treatment-naïve patients with WM who carry




FIG 2. Genomic-based treatment algorithm for symptomatic, treatment-naïve patients with Waldenström macroglobulinemia. Rituximab should be held for serum immunoglobulin M (IgM) $\geq 4,000$ mg/dL to prevent symptomatic IgM flare. Bendamustine and rituximab (Benda-R) should be considered as primary choice in patients with bulky adenopathy or extramedullary disease. Proteasome inhibitor (PI)-based regimens should be considered in patients with symptomatic amyloidosis, with autologous stem-cell transplantation as possible consolidation. Rituximab alone or with ibrutinib if *MYD88* is mutated or bendamustine may be considered in patients with IgM peripheral neuropathy (PN) depending on severity and pace of progression. Cyclophosphamide-based therapy such as dexamethasone, cyclophosphamide, and rituximab (DRC) represents an alternative to Benda-R but may be less effective. Maintenance rituximab may be considered in patients responding to rituximab-based regimens. BTK-I, Bruton tyrosine kinase inhibitor; CAGG, cold agglutininemia; CRYOS, cryoglobulinemia; HV, hyperviscosity; Mut, mutated; WT, wild type (not mutated).




FIG 3. Genomic-based treatment algorithm for symptomatic, previously treated or refractory patients with Waldenström macroglobulinemia. Rituximab should be held for serum immunoglobulin M (IgM) $\geq 4,000$ mg/dL to prevent symptomatic IgM flare. Bendamustine and rituximab (Benda-R) should be considered as primary choice in patients with bulky adenopathy or extramedullary disease. Proteasome inhibitor (PI)-based regimens should be considered in patients with symptomatic amyloidosis, with autologous stem-cell transplantation (ASCT) as possible consolidation. Rituximab alone or with ibrutinib if *MYD88* is mutated or bendamustine may be considered in patients with IgM peripheral neuropathy (PN) depending on severity and pace of progression. Rituximab combined with cyclophosphamide and dexamethasone (DRC) represents an alternative to Benda-R, although it may be a potentially less effective option. Maintenance rituximab may be considered in patients responding to rituximab-based regimen. (^a) Nucleoside analogues (NA) should be avoided in younger patients and candidates for ASCT. ASCT may be considered in patients with multiple relapses and chemotherapy-sensitive disease. BTK-I, Bruton tyrosine kinase inhibitor; CAGG, cold agglutinin; CRYOS, cryoglobulinemia; HV, hyperviscosity; Mut, mutated; WT, wild-type (not mutated).

only the *MYD88* mutation. Current nonrandomized comparisons of data from ibrutinib monotherapy and combined ibrutinib and rituximab studies show little difference in patients only have *MYD88* mutation; hence, ibrutinib alone can be considered for patients without bulky disease and symptomatic amyloidosis and for whom no contraindication to ibrutinib exists.⁵³⁻⁵⁶

Alternatives to ibrutinib that can be considered include bendamustine and rituximab (Benda-R) or PI-based therapy such as bortezomib, dexamethasone, and rituximab (BDR); carfilzomib, rituximab, and dexamethasone (CaRD); or ixazomib, dexamethasone, and rituximab (IDR). Randomized and nonrandomized studies suggest that cyclophosphamide-based therapy, although active in WM, may not be as effective as Benda-R or PI-based therapy.^{65,70-72}

Because the role of ibrutinib is not well established in patients with WM with bulky disease and has been lackluster in symptomatic patients with light chain (LC) amyloidosis, alternatives should be sought for these patients.^{33,73} Benda-R is a reasonable choice for those with bulky disease, ideally limited to 4 cycles to minimize prolonged cytopenias and long-term adverse events.^{64,65,71} Dexamethasone, rituximab,

and cyclophosphamide remains an alternative if Benda-R is not available.⁷² A PI-based treatment such as weekly BDR to minimize treatment-related neuropathy or use of neuropathy-sparing PI regimens such as CaRD or IDR may be considered for those with symptomatic LC amyloidosis.^{60,61,68,69,74} In these patients, consolidation with autologous stem-cell transplantation (SCT) may be considered.⁷⁵ Finally, patients with WM with IgM demyelinating neuropathy require deep and sustained responses to minimize myelin-directed monoclonal IgM deposition. For such patients, rituximab alone can be considered if the neuropathy is mild, but for more moderate to severe cases or patients with more rapid progression, the use of ibrutinib alone or with rituximab, Benda-R, or a regimen using a neuropathy-sparing PI such as CaRD or IDR may be considered.^{33,60,61,68,69,76}

For patients with both *MYD88* and *CXCR4* mutations, the necessity to achieve a rapid response must be considered. Nonsense *CXCR4*-mutated patients in particular are at risk for presenting with symptomatic hyperviscosity requiring immediate disease control.⁴¹ Rituximab should be avoided in patients with WM with high IgM levels or symptomatic hyperviscosity until such time as the IgM has decreased to

< 4,000 mg/dL to avoid a symptomatic rituximab flare.^{68,69} For patients requiring more emergent responses, either bendamustine or a PI-based regimen may be considered. For patients without symptomatic hyperviscosity and emergent need for disease control, ibrutinib plus rituximab may be preferred versus ibrutinib alone because the median time to achieve a major response appears to be shorter with the combination (3 v 6–7 months, respectively).^{53,54,56}

For patients presenting with wild-type *MYD88* WM determined by AS-PCR, Sanger sequencing or NGS that examines the TIR exon coding region of the *MYD88* gene should be done to exclude non-L265P *MYD88* mutations because patients with non-L265P *MYD88* mutations show responses to ibrutinib.¹² For patients with true *MYD88* wild-type WM, either bendamustine or PI-based therapy with rituximab can be considered because responses have been observed with these regimens in patients with both *MYD88* wild-type and mutated WM.^{60,64,65} In all patients with WM showing a response to a fixed-duration rituximab-containing regimen, maintenance rituximab may be considered.^{68,69} In patients who develop rituximab intolerance, ofatumumab may represent an option.^{77,78} There are no data at this time on the use of obinotuzumab in WM.

In previously treated patients with WM (Fig 3), the reuse of a front-line regimen may be considered for patients with long responses (> 2–3 years), although alternatives to alkylators or nucleoside analogs should be considered if previously used to minimize stem-cell damage. Nucleoside analogs should also be avoided in younger patients who may be candidates for autologous SCT and to minimize risk of disease transformation.⁶⁹ The use of ibrutinib can be considered in symptomatic, previously treated patients with mutated *MYD88* WM following the algorithm described earlier for symptomatic treatment-naïve patients with WM (Fig 2). For patients who have been exposed to ibrutinib, either Benda-R or a PI-based regimen may be considered. Alternatively, off-label use of acalabrutinib may be considered for patients not tolerating ibrutinib given findings in patients with chronic lymphocytic leukemia showing a successful switchover from ibrutinib for intolerant patients.⁷⁹ The findings of a prospective randomized trial of zanubrutinib versus ibrutinib (ClinicalTrials.gov identifier [NCT03053440](#)) may also provide important guidance on the use of zanubrutinib in patients not tolerating ibrutinib.

For patients who have experienced multiple relapses and have been previously exposed to alkylators, PIs, and ibrutinib, the off-label use of venetoclax may be considered. High overall response rates and durable responses were observed in patients with WM on venetoclax, including those who were *CXCR4* mutated and previously exposed to a BTK inhibitor, although response depth was affected by these factors.^{59,80} Everolimus, alone or in combination with bortezomib and rituximab, can also be considered in patients previously exposed to alkylators, PIs, and ibrutinib, although hematologic toxicities and autoimmune pneumonitis are noteworthy.^{63,81,82} Diminished activity in patients with wild-type *MYD88* or

mutated *CXCR4* has also been reported with everolimus and should be considered in context with other available treatments.⁶³ The use of cyclophosphamide, bortezomib, and dexamethasone, alone or with rituximab, is also an active regimen in previously treated patients with WM.⁸²

The use of autologous SCT can be considered in patients with multiple relapses and who remain chemotherapy sensitive.^{69,83} As mentioned earlier, autologous SCT may also be considered as a consolidative measure in patients with symptomatic amyloidosis.⁷⁵ Data on the efficacy and safety of allogeneic SCT or cellular therapies are limited and should ideally be considered in the context of a clinical trial. Finally, both treatment-naïve and previously treated patients should actively be considered for clinical trials. A listing of clinical trials for WM can be found at ClinicalTrials.gov.

MOLECULAR DIAGNOSTIC APPROACHES AND LIMITATIONS

In patients with suspected WM, the use of AS-PCR on BM aspirates represents the most optimal initial assay for detection of *MYD88* L265P mutations. If the patient is determined to have wild-type *MYD88* by AS-PCR, Sanger sequencing or targeted NGS that examines the TIR exon coding region of *MYD88* can be performed to evaluate for non-L265P *MYD88* mutations.¹⁰ Targeted NGS can also be used to identify *MYD88* L265P mutations, although false-negative results were observed in 30% of patients in one study when compared with AS-PCR, particularly in patients with low BM disease involvement.⁸⁴ The use of CD19-selected cells can improve chances of detection of mutated *MYD88*, although tumor selection is not routinely performed in clinical laboratories. CD19-selected PB mononuclear cells may also be used for AS-PCR testing of mutated *MYD88*, although diagnostic yield is greatly reduced with prior therapy, particularly rituximab, and use of unselected PB mononuclear cells.¹⁵

Targeted NGS or Sanger sequencing represents the best approach for evaluation of C-terminal mutations in *CXCR4* because > 40 nonsense and frameshift variants have been reported in WM.^{8,13,39,42} Because *CXCR4* mutations are subclonal with a median clonal distribution of 35%, false-negative results can occur in patients with low BM disease burden and with use of unselected mononuclear cells for *CXCR4* mutation testing.³⁹ Ultra-deep NGS (> 1,000×) may improve detection of *CXCR4* mutations over Sanger sequencing in patients with low tumor burden and clonality.⁸⁵ However, although large insertions or deletions in *CXCR4* are uncommon in WM, these may be missed with NGS but identified by Sanger sequencing.

SUMMARY

In summary, NGS has revealed recurring somatic mutations in WM that include *MYD88* and *CXCR4*. The genomic findings have contributed to ongoing targeted drug development and the recognition of a precision-guided treatment approach to WM. *MYD88* and *CXCR4* mutation status can be used to guide treatment decisions in WM.

AFFILIATIONS

¹Bing Center for Waldenström's Macroglobulinemia, Dana-Farber Cancer Institute, Boston, MA

²Department of Medicine, Harvard Medical School, Boston, MA

³Center for Prevention of Progression of Blood Cancers, Dana-Farber Cancer Institute, Boston, MA

⁴Clinical Research Development and Phase I Unit, CREA Laboratory, Azienda Socio Sanitaria Territoriale degli Spedali Civili di Brescia, Brescia, Italy

⁵Division of Hematology and Oncology, Massachusetts General Hospital, Boston, MA

CORRESPONDING AUTHOR

Steven P. Treon, MD, PhD, Bing Center for Waldenström's Macroglobulinemia, Dana-Farber Cancer Institute, M547, 450 Brookline Ave, Boston, MA 02215; e-mail: steven_treon@dfci.harvard.edu.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST AND DATA AVAILABILITY STATEMENT

Disclosures provided by the authors and data availability statement (if applicable) are available with this article at DOI <https://doi.org/10.1200/JCO.19.02314>.

AUTHOR CONTRIBUTIONS

Conception and design: Steven P. Treon, Mark Bustoros, Antonio Sacco, Aldo Roccaro, Irene M. Ghobrial

Administrative support: Christopher J. Patterson

Collection and assembly of data: Steven P. Treon, Lian Xu, Xia Liu, Maria Demos, Joshua Gustine, Mani Munshi, Nicholas Tsakmaklis, Jiaji G. Chen, Amanda Kofides, Andrew Keezer, Kirsten Meid, Christopher J. Patterson, Guang Yang, Irene M. Ghobrial

Data analysis and interpretation: Steven P. Treon, Maria Luisa Guerrero, Cristina Jimenez, Zachary R. Hunter, Gloria Chan, Romanos Sklaventis-Pistofidis, Andrew R. Branagan, Irene M. Ghobrial, Jorge J. Castillo

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

REFERENCES

1. Swerdlow SH, Kuzo I, Dogan A, et al: The many faces of small B cell lymphomas with plasmacytic differentiation and the contribution of MYD88 testing. *Virchows Arch* 468:259-275, 2016
2. Treon SP: How I treat Waldenström macroglobulinemia. *Blood* 114:2375-2385, 2009
3. Hunter ZR, Yang G, Xu L, et al: Genomics, signaling, and treatment of Waldenström macroglobulinemia. *J Clin Oncol* 35:994-1001, 2017
4. Treon SP, Gustine J, Xu L, et al: MYD88 wild-type Waldenström macroglobulinemia: Differential diagnosis, risk of histological transformation, and overall survival. *Br J Haematol* 180:374-380, 2018
5. Hunter ZR, Xu L, Tsakmaklis N, et al: Insights into the genomic landscape of MYD88 wild-type Waldenström macroglobulinemia. *Blood Adv* 2:2937-2946, 2018
6. Treon SP, Xu L, Yang G, et al: MYD88 L265P somatic mutation in Waldenström's macroglobulinemia. *N Engl J Med* 367:826-833, 2012
7. Jiménez C, Sebastián E, Chillón MC, et al: MYD88 L265P is a marker highly characteristic of, but not restricted to, Waldenström's macroglobulinemia. *Leukemia* 27:1722-1728, 2013
8. Poulain S, Roumier C, Decambron A, et al: MYD88 L265P mutation in Waldenström macroglobulinemia. *Blood* 121:4504-4511, 2013
9. Varettoni M, Arcaini L, Zibellini S, et al: Prevalence and clinical significance of the MYD88 (L265P) somatic mutation in Waldenström's macroglobulinemia and related lymphoid neoplasms. *Blood* 121:2522-2528, 2013
10. Xu L, Hunter ZR, Yang G, et al: MYD88 L265P in Waldenström macroglobulinemia, immunoglobulin M monoclonal gammopathy, and other B-cell lymphoproliferative disorders using conventional and quantitative allele-specific polymerase chain reaction. *Blood* 121:2051-2058, 2013
11. Gustine J, Meid K, Xu L, et al: To select or not to select? The role of B-cell selection in determining the MYD88 mutation status in Waldenström macroglobulinemia. *Br J Haematol* 176:822-824, 2017
12. Treon SP, Xu L, Hunter Z: MYD88 mutations and response to ibrutinib in Waldenström's macroglobulinemia. *N Engl J Med* 373:584-586, 2015
13. Varettoni M, Zibellini S, Defrancesco I, et al: Pattern of somatic mutations in patients with Waldenström macroglobulinemia or IgM monoclonal gammopathy of undetermined significance. *Haematologica* 102:2077-2085, 2017
14. Varettoni M, Zibellini S, Arcaini L, et al: MYD88 (L265P) mutation is an independent risk factor for progression in patients with IgM monoclonal gammopathy of undetermined significance. *Blood* 122:2284-2285, 2013
15. Xu L, Hunter ZR, Yang G, et al: Detection of MYD88 L265P in peripheral blood of patients with Waldenström's macroglobulinemia and IgM monoclonal gammopathy of undetermined significance. *Leukemia* 28:1698-1704, 2014
16. Alegría-Landa V, Prieto-Torres L, Santonja C, et al: MYD88 L265P mutation in cutaneous involvement by Waldenström macroglobulinemia. *J Cutan Pathol* 44:625-631, 2017
17. Poulain S, Boyle EM, Roumier C, et al: MYD88 L265P mutation contributes to the diagnosis of Bing Neel syndrome. *Br J Haematol* 167:506-513, 2014
18. Gustine JN, Meid K, Hunter ZR, et al: MYD88 mutations can be used to identify malignant pleural effusions in Waldenström macroglobulinemia. *Br J Haematol* 180:578-581, 2018
19. Drandi D, Genuardi E, Dogliotti I, et al: Highly sensitive MYD88^{L265P} mutation detection by droplet digital polymerase chain reaction in Waldenström macroglobulinemia. *Haematologica* 103:1029-1037, 2018
20. Bagratuni T, Ntanas-Stathopoulos I, Gavriatopoulou M, et al: Detection of MYD88 and CXCR4 mutations in cell-free DNA of patients with IgM monoclonal gammopathies. *Leukemia* 32:2617-2625, 2018
21. Hunter ZR, Xu L, Yang G, et al: The genomic landscape of Waldenström macroglobulinemia is characterized by highly recurring MYD88 and WHIM-like CXCR4 mutations, and small somatic deletions associated with B-cell lymphomagenesis. *Blood* 123:1637-1646, 2014
22. Tsakmaklis N, Xu L, Manning R, et al: Mutated MYD88 homozygosity is increased in previously treated patients with Waldenström's macroglobulinemia, and associates with CXCR4 mutation status and ibrutinib exposure. *Blood* 130, 2017 (abstr 1462)
23. Treon SP, Tsakmaklis N, Meid K, et al: Mutated MYD88 zygosity and CXCR4 mutation status are important determinants of ibrutinib response and progression free survival in Waldenström's macroglobulinemia. *Blood* 128, 2016 (abstr 2984)
24. Bustoros M, Sklaventis-Pistofidis R, Kapoor P, et al: Progression risk stratification of asymptomatic Waldenström macroglobulinemia. *J Clin Oncol* 37:1403-1411, 2019

25. Abeykoon JP, Paludo J, King RL, et al: MYD88 mutation status does not impact overall survival in Waldenström macroglobulinemia. *Am J Hematol* 93:187-194, 2018
26. Castillo JJ, Jurczynski A, Brozova L, et al: IgM myeloma: A multicenter retrospective study of 134 patients. *Am J Hematol* 92:746-751, 2017
27. Avet-Loiseau H, Garand R, Lodé L, et al: Translocation t(11;14)(q13;q32) is the hallmark of IgM, IgE, and nonsecretory multiple myeloma variants. *Blood* 101:1570-1571, 2003
28. Willenbacher W, Willenbacher E, Brunner A, et al: Improved accuracy of discrimination between IgM multiple myeloma and Waldenström macroglobulinemia by testing for MYD88 L265P mutations. *Br J Haematol* 161:902-904, 2013
29. Lin SC, Lo YC, Wu H: Helical assembly in the MyD88-IRAK4-IRAK2 complex in TLR/IL-1R signalling. *Nature* 465:885-890, 2010
30. Ngo VN, Young RM, Schmitz R, et al: Oncogenically active MYD88 mutations in human lymphoma. *Nature* 470:115-119, 2011
31. Yang G, Zhou Y, Liu X, et al: A mutation in MYD88 (L265P) supports the survival of lymphoplasmacytic cells by activation of Bruton tyrosine kinase in Waldenström macroglobulinemia. *Blood* 122:1222-1232, 2013
32. Yang G, Buhrlage SJ, Tan L, et al: HCK is a survival determinant transactivated by mutated MYD88, and a direct target of ibrutinib. *Blood* 127:3237-3252, 2016
33. Treon SP, Tripsas CK, Meid K, et al: Ibrutinib in previously treated Waldenström's macroglobulinemia. *N Engl J Med* 372:1430-1440, 2015
34. Phelan JD, Young RM, Webster DE, et al: A multiprotein supercomplex controlling oncogenic signalling in lymphoma. *Nature* 560:387-391, 2018
35. Paiva B, Corchete LA, Vidriales MB, et al: The cellular origin and malignant transformation of Waldenström macroglobulinemia. *Blood* 125:2370-2380, 2015
36. Argyropoulos KV, Vogel R, Ziegler C, et al: Clonal B cells in Waldenström's macroglobulinemia exhibit functional features of chronic active B-cell receptor signaling. *Leukemia* 30:1116-1125, 2016
37. Munshi M, Chen JG, Liu X, et al: SYK is activated by mutated MYD88 and drives pro-survival signaling in MYD88 driven B-cell lymphomas. *Blood Cancer J* 10:Article 12, 2020
38. Manček-Keber M, Lainšček D, Benčina M, et al: Extracellular vesicle-mediated transfer of constitutively active MyD88^{L265P} engages MyD88^{wt} and activates signaling. *Blood* 131:1720-1729, 2018
39. Xu L, Hunter ZR, Tsakmaklis N, et al: Clonal architecture of CXCR4 WHIM-like mutations in Waldenström macroglobulinemia. *Br J Haematol* 172:735-744, 2016
40. Hernandez PA, Gorlin RJ, Lukens JN, et al: Mutations in the chemokine receptor gene CXCR4 are associated with WHIM syndrome, a combined immunodeficiency disease. *Nat Genet* 34:70-74, 2003
41. Treon SP, Cao Y, Xu L, et al: Somatic mutations in MYD88 and CXCR4 are determinants of clinical presentation and overall survival in Waldenström macroglobulinemia. *Blood* 123:2791-2796, 2014
42. Poulain S, Roumier C, Venet-Cailault A, et al: Genomic landscape of CXCR4 mutations in Waldenström macroglobulinemia. *Clin Cancer Res* 22:1480-1488, 2016
43. Roccaro AM, Sacco A, Jimenez C, et al: C1013G/CXCR4 acts as a driver mutation of tumor progression and modulator of drug resistance in lymphoplasmacytic lymphoma. *Blood* 123:4120-4131, 2014
44. Cao Y, Hunter ZR, Liu X, et al: The WHIM-like CXCR4(S338X) somatic mutation activates AKT and ERK, and promotes resistance to ibrutinib and other agents used in the treatment of Waldenström's Macroglobulinemia. *Leukemia* 29:169-176, 2015
45. Cao Y, Hunter ZR, Liu X, et al: CXCR4 WHIM-like frameshift and nonsense mutations promote ibrutinib resistance but do not supplant MYD88(L265P)-directed survival signalling in Waldenström macroglobulinemia cells. *Br J Haematol* 168:701-707, 2015
46. Guerrero ML, Tsakmaklis N, Xu L, et al: MYD88 mutated and wild-type Waldenström's Macroglobulinemia: Characterization of chromosome 6q gene losses and their mutual exclusivity with mutations in CXCR4. *Haematologica* 103:e408-e411, 2018
47. Castillo JJ, Gustine JN, Meid K, et al: Low levels of von Willebrand markers associate with high serum IgM levels and improve with response to therapy, in patients with Waldenström macroglobulinemia. *Br J Haematol* 184:1011-1014, 2019
48. Davis RE, Ngo VN, Lenz G, et al: Chronic active B-cell-receptor signalling in diffuse large B-cell lymphoma. *Nature* 463:88-92, 2010
49. Nishizumi H, Horikawa K, Mlinaric-Rascan I, et al: A double-edged kinase Lyn: A positive and negative regulator for antigen receptor-mediated signals. *J Exp Med* 187:1343-1348, 1998
50. Jiménez C, Alonso-Álvarez S, Alcoceba M, et al: From Waldenström's macroglobulinemia to aggressive diffuse large B-cell lymphoma: A whole-exome analysis of abnormalities leading to transformation. *Blood Cancer J* 7:e591, 2017
51. Poulain S, Roumier C, Bertrand E, et al: TP53 mutation and its prognostic significance in Waldenström's macroglobulinemia. *Clin Cancer Res* 23:6325-6335, 2017
52. Gustine JN, Tsakmaklis N, Demos MG, et al: TP53 mutations are associated with mutated MYD88 and CXCR4, and confer an adverse outcome in Waldenström macroglobulinemia. *Br J Haematol* 184:242-245, 2019
53. Treon SP, Meid K, Gustine J, et al: Ibrutinib monotherapy produces long-term disease control in previously treated Waldenström's macroglobulinemia: Final report of the Pivotal trial (NCT01614821). *Hematol Oncol* 37:184-185, 2019 (suppl; abstr 135)
54. Treon SP, Gustine J, Meid K, et al: Ibrutinib monotherapy in symptomatic, treatment-naïve patients with Waldenström macroglobulinemia. *J Clin Oncol* 36:2755-2761, 2018
55. Dimopoulos MA, Trotman J, Tedeschi A, et al: Ibrutinib for patients with rituximab-refractory Waldenström's macroglobulinemia (iNNOVATE): An open-label substudy of an international, multicentre, phase 3 trial. *Lancet Oncol* 18:241-250, 2017
56. Buske C, Tedeschi A, Trotman J, et al: Ibrutinib treatment in Waldenström's macroglobulinemia: Follow-up efficacy and safety from the iNNOVATE study. *Blood* 132, 2018 (abstr 149)
57. Xu L, Tsakmaklis N, Yang G, et al: Acquired mutations associated with ibrutinib resistance in Waldenström macroglobulinemia. *Blood* 129:2519-2525, 2017
58. Trotman J, Tam CS, Mariton P, et al: Improved depth of response with increased follow-up for patients with Waldenström macroglobulinemia treated with Bruton's tyrosine kinase inhibitor zanubrutinib. *Proceedings of the 23rd Congress of the European Hematology Association, Stockholm, Sweden, June 14-17, 2018* (abstr PS1186)
59. Castillo J, Gustine J, Meid K, et al: Prospective Phase II study of venetoclax in patients with previously treated Waldenström's Macroglobulinemia. *Proceedings of the 23rd Congress of the European Hematology Association, Stockholm, Sweden, June 14-17, 2018* (abstr S854)
60. Castillo JJ, Meid K, Gustine JN, et al: Prospective clinical trial of ixazomib, dexamethasone, and rituximab as primary therapy in Waldenström macroglobulinemia. *Clin Cancer Res* 24:3247-3252, 2018
61. Treon SP, Tripsas CK, Meid K, et al: Carfilzomib, rituximab, and dexamethasone (CaRD) treatment offers a neuropathy-sparing approach for treating Waldenström's macroglobulinemia. *Blood* 124:503-510, 2014
62. Sklavenitis-Pistofidis R, Capelletti M, Liu CJ, et al: Bortezomib overcomes the negative impact of CXCR4 mutations on survival of Waldenström macroglobulinemia patients. *Blood* 132:2608-2612, 2018

63. Treon SP, Meid K, Tripsas C, et al: Prospective, multicenter clinical trial of everolimus as primary therapy in Waldenström macroglobulinemia (WMCTG 09-214). *Clin Cancer Res* 23:2400-2404, 2017
64. Laribi K, Poulain S, Willems L, et al: Bendamustine plus rituximab in newly-diagnosed Waldenström macroglobulinaemia patients: A study on behalf of the French Innovative Leukaemia Organization (FILO). *Br J Haematol* 186:146-149, 2019
65. Paludo J, Abeykoon JP, Shreders A, et al: Bendamustine and rituximab (BR) versus dexamethasone, rituximab, and cyclophosphamide (DRC) in patients with Waldenström macroglobulinemia. *Ann Hematol* 97:1417-1425, 2018
66. Castillo J, Gustine J, Xu L, et al: CXCR4 mutation subtypes impact response and survival outcomes in patients with Waldenström macroglobulinemia treated with ibrutinib. *Br J Haematol* 187:356-363, 2019
67. Gustine JN, Tsakmaklis N, Demos MG, et al: CXCR4 S338X clonality is an important determinant of ibrutinib outcomes in patients with Waldenström macroglobulinemia. Proceedings of the 10th International Workshop on Waldenström's Macroglobulinemia, New York, NY, October 11-13, 2018 (abstr W35)
68. Treon SP: How I treat Waldenström macroglobulinemia. *Blood* 126:721-732, 2015
69. Leblond V, Kastiris E, Advani R, et al: Treatment recommendations from the Eighth International Workshop on Waldenström's Macroglobulinemia. *Blood* 128:1321-1328, 2016
70. Rummel MJ, Niederle N, Maschmeyer G, et al: Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: An open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 381:1203-1210, 2013
71. Castillo JJ, Gustine JN, Meid K, et al: Response and survival for primary therapy combination regimens and maintenance rituximab in Waldenström macroglobulinaemia. *Br J Haematol* 181:77-85, 2018
72. Kastiris E, Gavriatopoulou M, Kyrtsonis MC, et al: Dexamethasone, rituximab, and cyclophosphamide as primary treatment of Waldenström macroglobulinemia: Final analysis of a phase 2 study. *Blood* 126:1392-1394, 2015
73. Pika T, Hegenbart U, Flodrova P, et al: First report of ibrutinib in IgM-related amyloidosis: Few responses, poor tolerability, and short survival. *Blood* 131:368-371, 2018
74. Gavriatopoulou M, García-Sanz R, Kastiris E, et al: BDR in newly diagnosed patients with WM: Final analysis of a phase 2 study after a minimum follow-up of 6 years. *Blood* 129:456-459, 2017
75. Sidiqi MH, Buadi FK, Dispenzieri A, et al: Autologous stem cell transplant for IgM-associated amyloid light-chain amyloidosis. *Biol Blood Marrow Transplant* 25:e108-e111, 2019
76. Dimopoulos MA, Tedeschi A, Trotman J, et al: Phase 3 trial of ibrutinib plus rituximab in Waldenström's macroglobulinemia. *N Engl J Med* 378:2399-2410, 2018
77. Castillo JJ, Kanan S, Meid K, et al: Rituximab intolerance in patients with Waldenström macroglobulinaemia. *Br J Haematol* 174:645-648, 2016
78. Furman RR, Eradat HA, DiRienzo CG, et al: Once-weekly ofatumumab in untreated or relapsed Waldenström's macroglobulinaemia: An open-label, single-arm, phase 2 study. *Lancet Haematol* 4:e24-e34, 2017
79. Awan FT, Schuh A, Brown JR, et al: Acalabrutinib monotherapy in patients with chronic lymphocytic leukemia who are intolerant to ibrutinib. *Blood Adv* 3:1553-1562, 2019
80. Davids MS, Roberts AW, Seymour JF, et al: Phase I first-in-human study of venetoclax in patients with relapsed or refractory non-Hodgkin lymphoma. *J Clin Oncol* 35:826-833, 2017
81. Ghobrial IM, Redd R, Armand P, et al: Phase I/II trial of everolimus in combination with bortezomib and rituximab (RVR) in relapsed/refractory Waldenström macroglobulinemia. *Leukemia* 29:2338-2346, 2015
82. Ghobrial IM, Witzig TE, Gertz M, et al: Long-term results of the phase II trial of the oral mTOR inhibitor everolimus (RAD001) in relapsed or refractory Waldenström Macroglobulinemia. *Am J Hematol* 89:237-242, 2014
83. Leblebjian H, Noonan K, Paba-Prada C, et al: Cyclophosphamide, bortezomib, and dexamethasone combination in Waldenström macroglobulinemia. *Am J Hematol* 90:E122-E123, 2015
84. Kyriakou C, Canals C, Sibon D, et al: High-dose therapy and autologous stem-cell transplantation in Waldenström macroglobulinemia: The Lymphoma Working Party of the European Group for Blood and Marrow Transplantation. *J Clin Oncol* 28:2227-2232, 2010
85. Kofides A, Demos M, Tsakmaklis N, et al: Alternative mutations and isoform dysregulation in MYD88 in Waldenström's macroglobulinemia. *Blood* 132, 2018 (abstr 1566)
86. Shin DW, Kim SM, Kim JA, et al: Characteristics of Waldenström macroglobulinemia in Korean patients according to mutational status of MYD88 and CXCR4: Analysis using ultra-deep sequencing. *Clin Lymphoma Myeloma Leuk* 19:e496-e505, 2019
87. Owen RG, Kyle RA, Stone MJ, et al: Response assessment in Waldenström macroglobulinaemia: Update from the VIth International Workshop. *Br J Haematol* 160:171-176, 2013



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**Genomic Landscape of Waldenström Macroglobulinemia and Its Impact on Treatment Strategies**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/journal/jco/site/ifc.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Steven P. Treon

Consulting or Advisory Role: Janssen, Pharmacyclics, BeiGene
Research Funding: Janssen, Pharmacyclics, Bristol-Myers Squibb
Travel, Accommodations, Expenses: Janssen Oncology
Other Relationship: Janssen, Pharmacyclics

Zachary R. Hunter

Honoraria: Janssen

Mark Bustoros

Honoraria: Takeda, DAVA Oncology
Consulting or Advisory Role: Takeda

Antonio Sacco

Consulting or Advisory Role: Celgene (I), Janssen-Cilag (I)
Research Funding: AstraZeneca (I)
Patents, Royalties, Other Intellectual Property: Pending patent: Treatment of C1013G/CXCR4-Associated Waldenström's Macroglobulinemia With an Anti-CXCR4 Antibody; application No.: 16/271569-BMS Ref. 12242-US-PCD
Travel, Accommodations, Expenses: Janssen-Cilag (I)

Aldo Roccaro

Consulting or Advisory Role: Celgene, Janssen-Cilag
Research Funding: AstraZeneca
Patents, Royalties, Other Intellectual Property: Pending patent: Treatment of C1013G/CXCR4-Associated Waldenström's Macroglobulinemia With an Anti-CXCR4 Antibody; application No.: 16/271569-BMS Ref. 12242-US-PCD
Travel, Accommodations, Expenses: Janssen-Cilag

Andrew R. Branagan

Consulting or Advisory Role: Surface Oncology, Pharmacyclics/Janssen

Guang Yang

Employment: AbbVie (I)
Honoraria: Janssen

Irene M. Ghobrial

Honoraria: Celgene, Bristol-Myers Squibb, Takeda, Amgen, Janssen, Karyopharm Therapeutics, Cellectar, Adaptive Biotechnologies, Sanofi, Medscape
Consulting or Advisory Role: Bristol-Myers Squibb, Novartis, Amgen, Takeda, Noxxon Pharma, Celgene, Sanofi, Genentech, GlaxoSmithKline, GNS Healthcare, Karyopharm Therapeutics, Adaptive Biotechnologies, Janssen, Medscape, AbbVie
Travel, Accommodations, Expenses: Bristol-Myers Squibb, Novartis, Onyx, Millennium, Celgene, Takeda, Janssen Oncology

Jorge J. Castillo

Consulting or Advisory Role: Pharmacyclics, Janssen, Vical, Genentech
Research Funding: Pharmacyclics (Inst), AbbVie (Inst), Janssen (Inst), BeiGene (Inst), TG Therapeutics (Inst)

No other potential conflicts of interest were reported.